

2019 - IID - MD4043 - eksamen 1
Eksamensdato: 2019-05-29

1

Which cause of infertility is the most difficult to treat in a couple who apply for assisted reproduction at a Norwegian infertility clinic?

- A The woman's BMI >30 and she smokes 10 cigarettes a day
- B The man has previously been sterilised and an attempted vasectomy reversal has not been successful
- C The woman is 38 years old and has an s-AMH <1.1 pmol/L
- D The man is 55 years old and his ejaculate contains very few motile sperm

000015de624e00ca30

2

A 35-year old woman has had amenorrhea for the last 8 months. She uses no form of hormonal contraception, or other medicines. The last three months she has noticed secretion from her breasts. She has had some problems with her vision over the last month. What is the most probable diagnosis?

- A Anorexia
- B Pregnancy
- C Prolactinoma
- D Primary ovarian failure

000015de624e00ca30

3

A 29-year old woman is pregnant in week 19. Routine ultrasound reveals a 6 cm tumour in one of her ovaries which most probably represents a fibroma. What is the most common complication she can get?

- A Degeneration
- B Rupture
- C Bleeding
- D Torsion

000015de624e00ca30

4

You are the GP for a 52-year old woman who was recently admitted to the Gynaecology Dept for acute abdominal pain. You have received the hospital discharge notes which state that due to a large, torsioned ovarian tumour the patient underwent bilateral laparoscopic salpingo-oophorectomy. The histology results reveal a serous cystadenoma with no atypia in the left ovary. Cytology of the peritoneal fluid reveals mesothelial cells. The patient contacts you for the results and for further treatment plan. She is in good health and has no problems after the intervention. You inform her about the test results.

What is the most correct next step for you to do now?

- A Write a prescription for combined oestrogen/progestin medication
- B Agree that she contacts you if she has increasing symptoms of hormone insufficiency
- C Refer her for surgery for staging at the Gynaecology Cancer Dept. and mark the referral "ovarian cancer pathway"
- D Refer her for follow-up with the gynaecologist for transvaginal ultrasound in 3 months

000015de624e00ca30

5

A 20-year old woman attends your GP surgery because she wants a contraceptive. She has had a legal medical abortion twice while she was on the contraceptive pill, and she most definitely does not want an intrauterine device.

What is the best option you can recommend her?

- A Vaginal contraceptive ring
- B Contraceptive implant
- C Condom
- D Postcoital contraception (emergency contraception)

000015de624e00ca30

6

Histologic type is one of several prognostic factors in endometrial cancer. Which of the histologic types mentioned has the best prognosis?

- A** Carcinosarcoma (mixed Müllerian tumour)
 - B** Serous papillary adenocarcinoma
 - C** Clear cell adenocarcinoma
 - D** Endometrioid adenocarcinoma
-

000015def24e00ca30

7

A 43-year old woman has an appointment with you as her GP. She has previously given birth to two children. She complains that over the last months she has had increasingly irregular and heavy menstrual bleedings which means that she has to change her underwear twice a day. The patient does not have hot flushes.

Which investigations should you do, i.e. which examinations and which samples should you take?

- A** Perform a gynaecological examination with cytology, and take blood samples for FSH and LH (gonadotropins, follicle stimulating hormone and luteinising hormone)
 - B** Take blood samples: Hb (haemoglobin), Ferritin and tumour marker CA125. Then send her to a specialist in gynaecology.
 - C** Take blood samples for oestradiol, progesterone, FSH and LH (gonadotropins, follicle stimulating hormone and luteinising hormone)
 - D** Perform a gynaecological examination with cytology and pipelle (endometrial biopsy); take blood samples: Hb.
-

000015def24e00ca30

8

A 28-year old woman attends your GP practice because of bleeding during her pregnancy. She has taken two positive pregnancy tests at home. She says that she has a definite date for her last period, and using the "Snurra" you determine that she is pregnant in week 7+2. She is G2 P0. One previous miscarriage after 8 weeks' of amenorrhoea. The bleeding that occurred the previous evening was fresh and comparable to that on the first day of a period. Now it is only a brownish discharge. The bleeding was accompanied by menstruation-like pain. Now, as the bleeding has stopped she has a stinging pain in the right iliac fossa. She has never had a cytology test (PAP smear) taken from her cervix.

How would you investigate/treat this patient?

- A** Perform a gynaecological examination to assess the cervix, uterus and adnexae. Take a cytology sample from the uterine cervix. Refer the patient to a gynaecologist for ultrasound querying an ectopic pregnancy.
 - B** Do not perform a gynaecological examination because she has just had a bleeding, and there is anyway no point in taking a cervical smear because the blood will interfere with the sample. Taking a sample will also provoke a new bleeding. Refer her directly to the Gynaecology department for ultrasound and assessment of the cervix since she is 28 years old and has never had a cervical cytology test (PAP smear).
 - C** Measure the blood concentration of HCG twice at a 3-7 day interval to determine whether the pregnancy is developing normally or not. Give her an appointment for a cervical cytology test (PAP smear) when the last HCG result is available.
 - D** Perform a gynaecological examination and listen for fetal heart beat (using a fetal doppler monitor for regular pregnancy check ups). If you can hear fetal heart beats to one of the sides, this could indicate an ectopic pregnancy and the patient should be referred to the Gynaecology department as an emergency.
-

000015def24e00ca30

9

A 35-year old woman gave birth to her second child 8 months ago. She is still somewhat breastfeeding. She has started back at work as a nurse and complains of large amounts of urinary incontinence. You find that she has a body mass index (BMI) of 36. Which treatment is the most correct for this patient now?

- A** Ask her to stop breastfeeding
 - B** Ask her to lose about 5 kg in weight
 - C** Recommend pelvic floor exercise with a physiotherapist
 - D** Recommend surgery for urinary incontinence (TVT, Tension-free Vaginal Tape procedure) in the Gynaecology Department
-

000015de624e00ca30

10

The patient is pregnant for the first time and has recently been for the routine ultrasound scan in week 19. The results revealed a placenta previa marginalis in which the lower edge of the placenta lay 3 mm from the internal cervical orifice. She has an appointment for a new placental examination in gestational week 32. You are her GP; at check-up with you she is anxious because she now has to have an elective Caesarian at term due to the position of the placenta. What can you tell her about the probability that the placental edge will have moved upwards away from the internal cervical orifice at the next ultrasound scan and a vaginal birth can be planned?

- A** 10%
 - B** 70%
 - C** 90%
 - D** 30%
-

000015de624e00ca30

11

What are the recommended procedures for treatment of genital herpes simplex virus (HSV) infection during pregnancy?

- A** An antiviral drug (aciclovir) is only suitable for pregnant women who are definitely known not to have had a previous genital HSV infection
 - B** If a pregnant woman gets a genital HSV infection during pregnancy, the fetus should be delivered by Caesarian section around the due date because of the increased risk of infection with a vaginal birth
 - C** Reactivation of genital HSV infection must always be treated with an antiviral drug (aciclovir)
 - D** Reactivation of genital HSV infection with frequent episodes towards the end of the pregnancy must be treated with an antiviral drug (aciclovir) 10 days before the due date
-

000015de624e00ca30

12

What is the most correct statement regarding listeria infection in pregnancy?

- A** Can result in serious complications in the fetus such as fetal death and neurological sequelae
 - B** Is tested for in everybody at the first pregnancy check-up
 - C** A common infection among pregnant women in Norway
 - D** Common symptoms are fever, nausea and abdominal pain
-

000015de624e00ca30

13

A 35-year old woman pregnant for the first time attends your GP surgery in gestational week 38+5. Previously she has generally been healthy and does not take any regular medication. The pregnancy has progressed normally and the patient has attended for regular check-ups with you. At today's check-up the patient complains about a frontal headache, but feels otherwise well. You measure her blood pressure to be 160/100, 158/97 and 165/95. Negative urine dipstick. What is the most correct recommendation for her?

- A** You say that since everything has been normal so far, she can take 1g paracetamol and see if her headache disappears
 - B** You recommend that she attends for a new blood pressure measurement in 5 days
 - C** You start her on antihypertensive medication, e.g. labetalol (Trandate®), and give her a new appointment in 5 days
 - D** You refer the patient to the Obstetrics Outpatient Clinic and ask for an assessment today or tomorrow
-

000015de624e00ca30

14

A young woman attends for an appointment with you, her GP. She gave birth to her 3rd child 2 months ago. She had diet-controlled gestational diabetes, otherwise it was an uncomplicated pregnancy and delivery.

What is the most correct follow-up for this woman?

- A** Ask her to keep a blood sugar diary and give her an appointment for follow-up in one week
 - B** Ask her to come back another day for a p-glucose test
 - C** Give her an appointment for an oral glucose tolerance test in a couple of weeks
 - D** Send a referral to the Endocrinology Outpatient Clinic at the nearest hospital
-

000015de624e00ca30

15

A 32-year old woman finds out that she is pregnant for the first time. She is 2 days over her expected period, the pregnancy is planned. When should she book an appointment with her GP for her first pregnancy check-up?

- A** In 1-2 weeks
 - B** In 8-12 weeks
 - C** In 4-8 weeks
 - D** In 2-4 weeks
-

000015de624e00ca30

16

You are a physician out in the districts. Travelling time to the hospital is three hours. A woman is gravida 3, para 2 with 2 normal pregnancies and deliveries. She attends for a planned check-up in gestational week 39 and you palpate a fetus in breech position.


What is the most correct information to tell her now?

- A** You believe that a breech delivery is associated with a high risk; you recommend she has a Caesarian section and refer her to the hospital tomorrow
 - B** You say you will refer her to the hospital for assessment of delivery method and possible external cephalic version
 - C** You say that you will perform an external cephalic version now, and ask the midwife in the office next door to assist you
 - D** You say that it is too late for an external cephalic version. This has to be done no later than gestational week 37
-

000015de624e00ca30

17

What does this ultrasound scan reveal? (The distance measured is 48 mm)



- A Cross-section of a fetal abdomen
- B A fetal head in the second trimester
- C A fetal head in the third trimester
- D A fetal head in the first trimester

000015def24e00ca30

18

Where does the fertilisation between an egg cell and a sperm cell usually take place?

- A In the ovary
- B In the cervix
- C In the Fallopian tube
- D In the uterine cavity

000015def24e00ca30

19

Ida (29 years old) is pregnant for the first time. She has checked out various websites for pregnant women and is feeling very confused by all the information she finds. She has a BMI of 35 and wonders if it is correct that she has an increased risk for many of the complications of pregnancy. Which of the complications below is Ida at increased risk of getting based on her BMI?

- A Breech delivery
- B Precipitate labour
- C Anaemia
- D Early spontaneous miscarriage

000015def24e00ca30

20

A 15-year old girl and her father attend the GP surgery for a medication check-up for the girl's ADHD. She was diagnosed with ADHD 1 year ago. Relevant treatment including medication was started in BUP. The girl takes methylphenidate 40 mg in the morning. You are told that over the last months the girl has had increasing problems at school with a lot of absences, she is sleeping a lot, has started self-harming, eats very little, does not take part in social activities or regular activities, and is irritable if she is not left alone. Which is the most probable comorbidity?

- A** Behaviour disorder
 - B** Personality disorder
 - C** Depression
 - D** Bipolar disorder
-

000015def24e00ca30

21

A 10-year old boy attends the doctor's surgery with his parents because the boy has increasingly made movements and sounds that bother him. He has blinked a little too hard and often for periods over the last few years. He has also had episodes of sniffing and throat clearing even though he doesn't have a cold. The last 6 months there has been a severe exacerbation with daily incidents with strong neck jerking and hitting and kicking the walls. Kicking the walls occurs mostly at home. In addition, his mother says that he can suddenly touch other people on the body, and that the parents of fellow pupils have contacted them to say they are shocked by his behaviour. He can breathe very heavily and exaggeratedly which has also scared others. The boy cries and wants to stop being bad. What would you do to help this patient?

- A** Refer to BUP for investigation and treatment
 - B** Refer the patient for MRI of the head and EEG to exclude a brain tumour and epilepsy
 - C** Treat the patient with Sertralin
 - D** Refer the family to the Family Counselling services for parental guidance
-

000015def24e00ca30

22

You are a doctor in general practice. A girl (14-years old) and her mother attend as an "emergency appointment". The girl has previously been healthy, amenable and done well at school. Over the last six months she has become more and more irritable, will not eat dinner with the family, and her mother hears her exercising in her room in the evenings. The mother has not seen her daughter vomit, but is concerned about a serious illness as the girl has lost several kilos in weight. The girl entered puberty around the age of 12, but has not had her periods the last 3 months. The patient says she is healthy, and that it's not a problem if she loses weight because she has thick thighs. At examination you find: Height 160cm, weight 36 kg (BMI 14.2 kg below the 2.5 percentile) BT 90/70, pulse 55, blood glucose 5, Hb 13. Apart from the loss of weight she appears to be in good health. Which further investigations and treatment do you think is most correct?

- A** Medical history, extended blood tests, mapping of diet and exercise patterns, and at the same time referral to the specialist healthcare services (BUP, Child and Adolescent Psychiatry Services)
 - B** Because the patient refuses to talk to you and she appears to be in good general health, you can't see any reason to waste time with an uncooperative patient, but encourage the mother to contact you if things get worse.
 - C** Because the weight loss has occurred gradually over several months and she appears to be in good general health, it is reasonable to agree an appointment in a week's time when the test results are back. You encourage her to increase her nutritional intake before the next appointment.
 - D** Eating disorders are a serious condition, and because the patient will not cooperate, voluntary admission with the parents' permission is recommended.
-

000015def24e00ca30

23

A mother comes to the Emergency Clinic with her 14-year old daughter after finding her crying in the bathroom with a rope she had tried to tie around her neck and some superficial cuts on her left wrist. The girl confirms that she had thoughts of wanting to die, and her mother is worried. The mother says that her daughter was admitted to a psychiatric day unit one year ago because of cuts to her left wrist. She then received outpatient follow-up for depression for a while, but stopped six months ago because she had improved. Over the last weeks she has been more and more off school and mostly stays in her room. The girl is quiet, looks down, makes little eye contact and will not say anything about how she feels or what she is thinking. She has some old scars from previous self-harming. What is your assessment of the suicide risk for this girl?

- A** I cannot assess the suicide risk before I have talked more with the girl about suicidal thoughts and suicidal plans.
 - B** I assess the suicide risk to be high, mostly due to the probable presence of a mental disorder, an increasing degree of withdrawal and confirmed suicidal thoughts.
 - C** I assess the suicide risk to be high, mostly due to the probable presence of a mental disorder, previous suicide attempt and confirmed suicidal thoughts.
 - D** I assess the suicide risk to be high mostly due to the probable presence of a mental disorder and repeated self-harming.
-

000015def24e00ca30

24

Why is it important to investigate personality disorder in adolescents with emotional difficulties, social relationships and self-harming over several years?

- A** To ensure the correct treatment
 - B** To have information on criteria ready for when the person is 18 years old.
 - C** Because a personality disorder indicates that there is no evidence-based treatment; the case should therefore be concluded in the specialist healthcare services and receive any follow-up in the form of support meetings in the first line services.
 - D** Because it will mean that she is a poor informant about her own health.
-

000015def24e00ca30

25

Why is it often more difficult to identify psychotic disorders in children and adolescents than in adults?

- A** Because the prevalence is much lower.
 - B** Because they are still developing and the symptoms are often less clear than in adults.
 - C** Because children and adolescents often will not talk about psychotic symptoms
 - D** Because children and adolescents have problems distinguishing between fantasy and reality.
-

000015def24e00ca30

26

A 5-year old boy has been diagnosed with conduct disorder.
What is the first line of treatment for this disorder?

- A** Systemic family therapy that improves the relationship between the parents so that the child, who is the symptom-bearer, stops his/her problematic behaviour.
 - B** Cognitive-oriented behavioural therapy with exposure and response prevention which targets the child's anxiety that underlies the behavioural problem
 - C** A development support guidance programme which promotes positive cooperation between the child and other children and between the child and adults.
 - D** Psychodynamic-oriented individual play therapy in which the child's inner conflicts are resolved so that the problematic behaviour stops
-

000015def24e00ca30

27

A 15-year old boy comes to the doctor's surgery together with his mother. He is pale, moves slowly, turns away and avoids answering questions. He has little facial expression. He has developed and functioned normally until recently. The last 5 weeks he has barely been to school and does not have contact with friends other than via the Internet. He doesn't talk and is sullen, and he can be very angry sometimes. She is worried and anxious because she feels she no longer has contact with him. He thinks things are fine, but is bored at school and thinks his mother is making a big fuss. He is awake all night, she doesn't know if he is playing games or surfing the Internet. He eats little, and is impossible to get up in the morning. Before he was active with ski jumping, but hasn't been to training the last 2 months because he doesn't think it is any fun anymore. What is the most probable diagnosis?

- A** Severe depressive episode with psychotic symptoms
 - B** Severe depressive episode without psychotic symptoms
 - C** Moderate depressive episode
 - D** Mild depressive episode
-

000015de624e00ca30

28

A 10-year old girl attends your GP surgery together with her father. For the last two weeks she has complained almost daily about a headache. She doesn't want to go to school in the morning, and is tired in the afternoons. She is fine the days she can be at home. The headache normally passes after 5-10 minutes. She has previously been healthy with normal development. Her parents are divorced, but cooperate well together. Her little sister who is 5 has recently been diagnosed with type 1 diabetes mellitus. Her best friend changed schools 3 months ago, and she has had problems fitting in socially since then. What would you do to help the patient?

- A** Provide psychoeducation, refer to a paediatrician for investigation and assessment for treatment
 - B** Provide psychoeducation, refer for MRI of the head, treat with paracetamol. New check-up in 2 weeks.
 - C** Provide psychoeducation, refer to BUP for investigation and assessment for treatment
 - D** Provide psychoeducation, important to get a good daily rhythm. Recommend adaptation at the school.
-

000015de624e00ca30

29

8-year old Siri's mother contacts you with the following concerns: During the last months, Siri has stopped eating sausages because she is afraid a piece could get stuck in her throat. She changes her socks several times a day because she is afraid she could have stepped in something nasty on the floor. When she gets home from school she changes all her clothes because she thinks her room gets dirty if she sits down with her school clothes on. If a friend visits she makes sure that she doesn't go into her room. The mother thinks she spends far too long in the shower. She mentions that it all started after they had had information about swine flu at school. What is the most probable explanation of her symptoms?

- A** Siri has generalised anxiety disorder.
 - B** Siri has paranoid thoughts and is developing a psychosis.
 - C** At this young age it is normal to react with anxiety to information about serious disease.
 - D** Siri has a compulsive-obsessive disorder
-

000015de624e00ca30

30

Aleksander is a 15-year old boy who in the autumn comes with his father to the GP surgery for assessment of tiredness and chronic stomach ache, which he localises down to the right fossa, which began two months ago around the time he started back at school. They were on holiday in Turkey early in August, but he had no symptoms of infection on holiday; he hasn't had a fever. He often sleeps when he comes home from school which he enjoys and where he has many friends. Father is the trainer for his football team and he is concerned that Aleksander is performing poorly. Aleksander does not have diarrhoea and says he has a daily bowel movement and that it is "normal". He says there is no blood or mucous. He has not vomited, but can feel nauseous. He is eating less and his father feels that he has become thinner over the last month. At examination you find he is slim, a little pale, sunken eyes, but otherwise in good general health. Has slight palpation tenderness low in the abdomen on the right side. No masses. Some inguinal glands. What is the most correct course of action that should be done next?

- A** Take a stool sample for infectious gastroenteritis.
 - B** Requisition US abdomen to exclude pathology.
 - C** Start treatment for obstipation; a new appointment in two months.
 - D** Take blood samples including gluten antibodies, hemofec and Calprotectin in faeces.
-

000015de624e00ca30

31

Mari is a 2-month old girl who is fully breastfed. She wakes very often and cries at night. The parents interpret this to be stomach pains. She sleeps well during the day for 2 hours. She eats well and is following her weight and length curve. She does not particularly gulp a lot. She swallows a lot during the night, and makes noises that her parents interpret as gulping that she swallows again. Her parents suspect that she has reflux (GER). At clinical examination in the surgery you find a healthy-looking child in normal good health without any specific findings. What do you do next?

- A** Start with conservative treatment with a high head position and more frequent small meals.
 - B** Explain to the parents about normal conditions.
 - C** Start treatment with a proton pump inhibitor (PPI) based on the clinical findings.
 - D** Refer her to the Paediatric Department for further investigations.
-

000015de624e00ca30

32

A 3-day old baby is to be discharged from the Maternity ward. Possible VSD was diagnosed in utero and after the birth the Paediatric Cardiologist confirms a moderately-large VSD. The family have been told they will have an Outpatient appointment in 4 weeks. The grandmother who works in the Cardiovascular Dept. asks if this is correct.

What do you think is the reason the doctor has given them this appointment?

- A** The relatively high lung pressure in the weeks after birth counteracts left to right shunting
 - B** VSD is associated with right to left shunting which rarely induces heart failure.
 - C** The size of the VSD will not increase until later
 - D** In neonates the heart does not work very hard because they are very inactive and sleep a lot in the first weeks.
-

000015de624e00ca30

33

A 2-year old previously healthy boy living in Norway has had a cold and fever for a day. The doctor at the Walk-in Centre ("legevakt") finds that he is lethargic, but not poorly, and that he has suppurative otitis media in one ear.

Which treatment would you recommend?

- A** Paracetamol dose x 3-4 until pain-free
 - B** Ibuprofen dose x 3-4 until pain-free
 - C** Phenoxyethylpenicillin for 5 days
 - D** Course of amoxicillin for 5 days
-

000015de624e00ca30

34

A 26-month old boy with known egg allergy. Has had all his vaccinations. Admitted with RSV bronchiolitis aged 6 months. Cold and fever initially for 1 week which has persisted with nightly cough and mucous production for the last 2 months. Is in nursery school where he is continually a little behind the other children in physical activities. At examination he has prolonged expiration and uses the abdominal help muscles. What is the most probable diagnosis?

- A** Whooping cough
- B** RSV bronchiolitis
- C** Pneumonia
- D** Asthma

000015d0e24e00ca30

35

A 7-month old boy. The father has allergy and eczema. Fully breastfed, the mother has no dietary restrictions, introduced to other food from 5 months of age. Subsequently he has tasted most of the family's food. Dry and itchy rash increasing over the last 2 weeks. What is the most probable diagnosis?



- A** Allergic urticaria
- B** Atopic dermatitis due to milk allergy.
- C** Atopic dermatitis probably not due to milk-protein allergy
- D** Chronic urticaria due to milk allergy.

000015d0e24e00ca30

36

You are a general practitioner. A mother attends with her 1-year old boy. The mother says that a short while ago a lump appeared in the boy's right groin. The lump is not always present, but it generally appears when the boy cries. When the mother squeezes the lump it is soft and can be pushed inwards. When you examine the boy in your office, he is quiet and you cannot find a lump. What is the correct action to take?

- A** You refer for ultrasound querying a swollen lymph node in the groin.
- B** You reassure the mother saying that you cannot find anything wrong when examining the groin.
- C** You suspect an inguinal hernia and refer to a Surgical Dept for an operation.

000015d0e24e00ca30

37

You are the doctor at the Child Health Centre. A 1-year old boy has only his left testicle in the scrotum. You think you can feel the right testicle in the right groin, but it is not possible to milk it down into the scrotum. What is the correct diagnosis?

- A Undescended testicle/testicle retention/retentio testis
- B Phimosis
- C Hydrocele
- D Inguinal hernia

000015de624e00ca30

38

You are the doctor in the minor injuries centre. A 13-year old boy has acute pain in the scrotum. The pain is severe and the boy is a little nauseous. What is the correct course of action?

- A Send the boy home and tell him to come back if the pain does not get better
- B Suspect testicular torsion and refer as emergency help to the Surgical Dept
- C Start antibiotic treatment for suspected epididymitis

000015de624e00ca30

39

A 10-year old girl has been on holiday in India. After coming home, she had watery diarrhoea that lasted more than 3 weeks. She has lost 1 kilo in weight, and complains of burping that tastes of rotten eggs. What is the most probable agent?

- A Giardia Lamblia
- B *Staphylococcus aureus*
- C Rotavirus
- D *Campylobacter*

000015de624e00ca30

40

CP is a collective name for a group of conditions characterised by a dysfunction of the movement and motor functions. The condition is caused by damage to the immature brain before the child is 2. What type of additional problems is a child with CP at increased risk of?

- A Impaired vision
- B All
- C Behaviour problems
- D Epilepsy
- E Cognitive problems

000015de624e00ca30

41

Petter is a 7-month old, previously healthy boy. He started with vomiting and diarrhoea six days ago. The last 24 hours, blood spots and mucous have appeared in his stools. He has also become more lethargic and drinks little. His mother takes him to the GP who finds he is pale, has a pulse 140/minute, temperature 37.2 degrees C, slightly cold peripherally, dry lips and slightly halonated eyes. Petter is referred to the hospital where investigations in the Emergency Department reveal the following:

Test	Result	Ref. range
CRP	27 mg/L	<5 mg/L
Leucocytes	16.9 $10^9/L$	3.7 – 14.7 $\times 10^9/L$
Hb	7.0 g/dL	10.5 – 13.1 g/dL
Thrombocytes	120 $\times 10^9/L$	229 – 497 $\times 10^9/L$
Creatinine	63 micromol/L	14 – 34 micromol/L

Which disease do you primarily suspect and which test best clarifies the diagnosis the same evening?

- A Acute lymphatic leukemia: blood smears, urate
- B Acute gastroenteritis: S/B, Na, K, faeces culture
- C Haemolytic uremic syndrome: LD, blood smear
- D Idiopathic thrombocytopenia: ANA, blood smear, hemo FEC

000015de624e00ca30

42

Children born with this condition often require two surgical procedures in the neonate period. One is to close the skin defect. What is the reason for the other surgical procedure?



- A Hydrocephalus
- B Club foot/Pes equinovarus
- C Tethered cord
- D Hip dysplasia

000015de624e00ca30

43

You are contacted by the district nurse as the GP for a 5-year old boy who has been for a check-up at the Child Health Centre that day. She noticed that the boy had poor dental hygiene and a lot of bad teeth; he also had a bruise on his left ear lobe that was painful. When she asked the boy how he got it he said that he "didn't remember" at the same time as he appeared upset and peeked up at his mother. How would you manage this and what advice would you give to the district nurse?

- A** You ask the district nurse to make sure the boy is referred to a dentist.
 - B** You give the boy an appointment for examination by you the same day, and consider a referral to the dentist and a note of concern to the Child Welfare Services.
 - C** You request an appointment for the boy in your surgery in 4 weeks.
 - D** You say you will call the mother to find out what happened.
-

000015def24e00ca30

44

Nils who is 8 years old has Diabetes Mellitus Type 1. He wakes up one morning feeling clammy and sweaty with a blood glucose of 11.0 mmol/L. How is this best explained?

- A** Night hypoglycemia with hormonal counterregulation in the form of increased excretion of glucagon and decreased excretion of adrenaline and growth hormone.
 - B** Night hypoglycemia with hormonal counterregulation in the form of increased excretion of adrenalin, glucagon and thyroxine.
 - C** Night hypoglycemia with hormonal counterregulation in the form of reduced excretion of adrenaline and increased excretion of glucagon and leptin.
 - D** Night hypoglycemia with hormonal counterregulation in the form of increased excretion of adrenaline, glucagon and growth hormone.
-

000015def24e00ca30

45

There are many reasons why a child can have an increased risk of delayed psychomotor development. Which of the following groups/answers are the most correct?

- A** Children who are born extremely premature
 - B** Children with a congenital heart disorder
 - C** All
 - D** Children with epilepsy
-

000015def24e00ca30

46

As a doctor you meet a 3-year old girl. Over the last weeks she has had bruises on her legs, and some small red spots have also been seen on her calfs. She is otherwise healthy now, but had a respiratory tract infection a few weeks ago. At clinical examination she is afebrile and in good general health. You find petechiae and bruises on her legs. Otherwise normal results at general clinical examination. General blood tests reveal thrombocytes $28 \times 109/L$ (ref. range: $228-435 \times 109/L$). All other relevant blood tests are normal. You are almost certain that the patient has ITP (Idiopathic thrombocytopenic purpura).

What is the correct way to manage this situation?

- A** Intravenous immunoglobulins
 - B** Oral steroids
 - C** Intravenous Rituximab
 - D** Continued observation
-

000015def24e00ca30

47

Which statement is correct?

- A** Iron deficiency is the most common cause of anaemia in children.
 - B** Iron supplements for children should be combined with milk to improve iron uptake in the intestines.
 - C** At the time of diagnosis of acute leukaemia in children, a hypochromic microcytic anaemia is often present.
 - D** In haemolytic uremic syndrome (HUS) in children, thrombocytopenia is often present.
-

000015def24e00ca30

48

As a specialty registrar, you are working at a GP's office where the first patient of the day is a 3-year old boy. The mother says that he has been down with a number of different respiratory tract infections over the last months and she thinks he is paler than normal. Clinical examination reveals a temperature of 38.4°C and poor general health. You find multiple small glands in his neck, enlarged tonsils and hepatosplenomegaly. Preliminary blood tests at the GP's office are presented in the Table below. What is the most probable diagnosis for this child?

Test:	Result:	Reference range:
CRP	36 mg/L	<5 mg/L
Hb	8.3 g/dL	10.5-13.1 g/dL
Thrombocytes	126 x 10 ⁹ /L	228-435 x 10 ⁹ /L
Leukocytes	15.9 x 10 ⁹ /L	3.7-14.7 x 10 ⁹ /L

- A Infectious mononucleosis
- B Hemolytic uremic syndrome
- C Acute leukaemia
- D Parvovirus B19

000015def24e00ca30

49

Kari is 5 years old, weight 20kg and has a nut allergy. At a birthday party where they served cake, she suddenly became red in the face, distant, and started gasping for breath. The emergency services doctor arrived quickly and is going to administer adrenaline. What dose should the doctor give?

- A Adrenaline 0.1 mg/ml at a dose of 2 ml i.v.
- B Adrenaline 0.01 mg/ml at a dose of 0.2 ml i.m.
- C Adrenaline 0.1 mg/ml at a dose of 0.2 ml i.m.
- D Adrenaline 1 mg/ml at a dose of 0.2 ml i.m.

000015def24e00ca30

50

A 65-year old man was admitted as an emergency. He was at a party and when dancing swing collided with another dancer. He got the other dancer's elbow in his left flank. He complains about severe pain in his left flank and has macroscopic haematuria. Findings at examination: BP 140/85 mmHg, Pulse 94/min. No visible injury at examination. Haemoglobin at admission is 11.6 (ref. range 13.4 - 17.0 g/dL.) Palpation tenderness in the left part of his abdomen and palpation tenderness above the costovertebral angle. CT of the kidneys was performed, see the images. What is the treatment?



- A** CT abdomen reveals hydronephrosis which should be treated with a JJ-stent.
- B** In the event of a fall in haemoglobin, the patient should undergo emergency surgery with a left-sided nephrectomy.
- C** Observation of the patient with check of HgB and vital parameters.
- D** Ultrasound-guided percutaneous drainage of the subcapsular haematoma.

000015de624e00ca30

51

72-year old man, previously generally healthy but with mild to moderate micturition problems. In connection with an operation for hip arthrosis, he was given a spinal anaesthetic without insertion of a bladder catheter. The operation was difficult and lasted a long time. Postoperatively the patient had problems with frequent, small volumes of urine and urine leakage. Urine dipstick was negative. What type of incontinence is the most probable in this patient?

- A** Stress incontinence
- B** Urge incontinence
- C** Mixed incontinence
- D** Overflow incontinence

000015de624e00ca30

52

A 76-year old man comes as an emergency help appointment to your surgery. Over the last two days he has had pain in the head of his penis with redness and swelling, and he cannot pull the prepuce forward. There is no swelling along the shaft of the penis. The pain and the swelling occurred the morning after he had "successful" unplanned intercourse in the town thanks to his GP having prescribed viagra tablets.

- A** He has been infected and has a bacterial balanitis.
- B** He has priapism secondary to an overdose of viagra.
- C** He has paraphimosis.
- D** He has urethritis probably with chlamydia.

000015de624e00ca30

53

A 33-year old man was re-admitted as an emergency 4 days after surgery for appendicitis. The operation was difficult with conversion from laparoscopy to open surgery due to bleeding and poor overview. At admission he complained of increasing pain in the right flank and nausea since the day of discharge after the operation. He has cold sweats, chills and is pale. Pulse: 124/min, blood pressure 80/50 and T_p 39.5°C. Blood tests reveal the following:

CRP	220 mg/L	0 - 5
B-HB	14.2 g/dL	13.4 - 17.0
B-Leucocytes	15.6 10 ⁹ /L	3.7 - 10.0
P-Sodium	145 mmol/L	137 - 145
P-Potassium	4.2 mmol/L	3.5 - 4.4
P-Creatinine	106 micromol/L	60 - 105

Emergency ultrasound of the abdomen revealed right side hydronephrosis and hydroureter as well as moderate amounts of free fluid in the pelvic cavity. The time is 23:00 in the evening. What is the appropriate treatment for this patient?

- A** The patient should go directly to the operating theatre for explorative laparotomy due to probable postoperative abscess.
- B** The patient should be given intravenous antibiotics and undergo emergency insertion of a nephrostomy.
- C** The patient should be given intravenous antibiotics and referred for ultrasound-guided drainage of abscess in the pelvic cavity.
- D** The patient should be given intravenous antibiotics and undergo emergency surgery with insertion of a JJ stent in the right ureter.

000015de624e00ca30

54

A woman aged 60 had macroscopic hematuria. Examination with cystoscopy showed a tumour in the urinary bladder. Transurethral resection of the bladder tumour (TURB) has been performed. Histological examination showed non-muscle-invasive urothelial cancer with small areas with *carcinoma in situ* (Ta GII w/ cis.). What is the best action?

- A** Radical cystectomy or radical radiation therapy
 - B** Immunotherapy (BCG) within four weeks, and check-up cystoscopy in three months
 - C** Check-up cystoscopy in three months
 - D** Re-TURB and if no tumour, check-up cystoscopy in three months
-

000015d0e24e00ca30

55

A 65-year old man with insulin-dependent diabetes mellitus and hypertension, well-regulated on antihypertensive drugs, is admitted as a semi-acute emergency to the Surgical Department with the diagnosis "tumour abdomen". He has had nightly urine leakage for the last 3 months. He is afebrile, creatinine is 250 µmol/L (reference: 60 - 105 µmol/L). CT abdomen taken the same evening shows a urinary bladder with upper/cranial edge between the umbilicus and xiphoid process, and bilateral hydronephrosis grade III. The urinary bladder contains 6,000 ml. What investigations and treatment should the doctor initiate?

- A** Insertion of a permanent catheter should immediately be done and surgery/TURP within 1 week if the creatinine normalises should be planned.
 - B** Insertion of a permanent catheter should immediately be done, and then ultrasound-guided bilateral nephrostomies as semi-acute help the next morning should be performed.
 - C** Clean intermittent catheterisation (CIC) should immediately be started, and requisition of cystometry/flowmetry, ultrasound of the kidneys and check of creatinine for the next day should be planned.
 - D** Insertion of a permanent catheter to ensure optimal drainage should be done immediately, and requisition of cystometry/flowmetry, blood tests, and new assessment at Urology Outpatients in 2 months should be planned.
-

000015d0e24e00ca30

56

A 20-year old man has an appointment at the health centre. One of his testicles has increased in size over the last 2-3 weeks. He has no pain. At examination, you feel that one of his testicles is uneven and hard compared to the healthy side, and you suspect testicular cancer. Otherwise you find nothing wrong at clinical examination. Which of the answers below is the best in regard to further investigation of the patient?

- A** You order US (ultrasound) of the scrotum, CT of the abdomen/thorax and take blood samples for tumour markers. You give a new appointment for information.
 - B** You explain to the patient that you suspect testicular cancer, and therefore refer the patient directly to the Urology Dept. for rush investigations without taking any samples.
 - C** You order blood tests with tumour markers and request a rush appointment, within 3 weeks for US (ultrasound) of the scrotum. You give the patient a new appointment after the ultrasound scan.
 - D** You explain to the patient that this could be a tumour, but that investigations with blood tests and ultrasound (US) within a week will be able to clarify this. You take blood samples with tumour markers. You order US of the scrotum and at the same time refer the patient to the Urology Dept. for testicular biopsy.
-

000015d0e24e00ca30

57

A young woman aged 20 goes to the doctor's surgery due to visible blood in her urine, diffuse pain in the lower abdomen and a burning sensation when urinating. You see the patient, examine her and find tenderness in the lower area of the abdomen at palpation. The urine sample is too bloody for the urine dipstick results to be reliable.

What do you do?

- A** You refer the patient for cystoscopy and 3-phase CT of the urinary tract.
- B** You start antibiotic treatment and order CT of the urinary tract without contrast so that the patient is not exposed to too much radiation.
- C** Because the patient has symptoms of a UTI you start treatment with antibiotics and ask her to come back in a week for a check-up, or earlier if things do not improve quickly.
- D** You send the urine to culture and order CT of the urinary tract. You will contact the patient when the results of the culture are ready so that you can prescribe the correct antibiotics.

000015de624e00ca30

58

Kristian, aged 72, is an active pensioner, and has been referred by his GP to Urology Outpatients due to recurring lower urinary tract infections over the last 18 months; the last time complicated with epididymitis. He has healthy heart and lungs, has insulin-requiring diabetes mellitus, and was operated 3 years ago for spinal stenosis (benign) at level L4-L5. Urination is characterised by hesitancy, reduced stream pressure, long voiding time. Periodic urine leakage, but no urge. Normal values for haemoglobin, creatinine and prostate specific antigen (PSA). Urine dipstick reveals white cells 3+ and nitrite 2+. Ultrasound urinary tract reveals no hydronephrosis, but 400 ml residual urine. Prostate volume is measured at 70 ml.

Which initial investigation and treatment is most suitable for this patient?

- A** Insert a permanent bladder catheter and refer the patient for TURP (transurethral resection of the prostate)
- B** Start with antibiotics for presumed cystitis and at the same time refer the patient for TURP within 2 weeks.
- C** Train Kristian to perform CIC 3-5 times a day (Clean Intermittent Catheterisation) and start on Duodart (alpha-blocker + 5-alpha-reductase inhibitor)
- D** Train Kristian to perform CIC 3-5 times a day and refer for investigation with cystoflowmetry (Pressure-flow investigation).

000015de624e00ca30

59

Harald (50 years old) has diabetes mellitus type II and uses Metformin. Over the last years he has experienced gradually increasing problems with impotence. He has an appointment with you his GP and wants a tablet that will help with his impotence problems.

Which statement is (most) correct?

- A** Viagra is a prostaglandin E1 inhibitor.
- B** PDE5 inhibitors are not contraindicated at concomitant use of nitroglycerine.
- C** There is only one type of phosphodiesterase type 5 (PDE5) inhibitor on the market.
- D** Viagra is a phosphodiesterase type 5 (PDE5) inhibitor.

000015de624e00ca30

60

In secondary hyperparathyroidism, PTH hormone production in the parathyroid gland is increased in response to an external stimulus, in contrast to primary hyperparathyroidism. A change in the blood concentration of phosphate and calcium can contribute to the increased PTH production.

Which of the alternatives will result in secondary hyperparathyroidism?

- A** Hypophosphatemia and hypercalcemia
- B** Hyperphosphatemia and hypocalcemia
- C** Hyperphosphatemia and hypercalcemia
- D** Hypophosphatemia and hypocalcemia

000015de624e00ca30

61

A 60-year-old woman is admitted after a three-day history with a cough and high fever. She is being treated for diabetes mellitus. On admission, she has stable vital parameters with blood pressure of 120/70 and pulse-oxymetry 94% without additional oxygen. However, her s-creatinine is 450 micromol/L (normal <90). Urine dipstick is negative for glucose, blood, proteins and leukocytes. U-sodium 10 mmol/L (low), u-osmolality 500 (medium to high).

What is the most likely cause of her acute kidney injury?

- A** Postrenal obstruction, and it should be considered whether the radiologist should perform a nephrostomy within a few hours.
- B** Acute tubular necrosis due to ischaemia/nephrotoxins
- C** Prerenal cause
- D** Acute glomerulonephritis (probably IgA nephritis caused by a throat infection)

000015def24e00ca30

62

A 28-year old man is referred to the Medical Outpatients Clinic because his GP has found slightly increased serum creatinine (115 umol/L, normal 60-105). The samples are otherwise normal apart from a markedly low s-bicarbonate (14 mmol/L, normal 22-28 mmol/L). He is otherwise healthy and fit. Which of the following renal diseases are characterised by early development of metabolic acidosis?

- A** Primary glomerulonephritis
- B** Kidney diseases in autoimmune system disorders
- C** Tubular kidney diseases
- D** Diabetic kidney disease

000015def24e00ca30

63

Medicines that affect the renin angiotensin system are used frequently and for many conditions. Which haemodynamic effects do these medicines have on renal circulation?

- A** The afferent arteriole is dilated strongly so that blood flow increases and renal function improves
- B** The efferent arteriole is dilated so that the GFR drops
- C** The afferent arteriole is dilated somewhat, but the efferent is dilated even more so that the capillary pressure is reduced considerably and the GFR drops
- D** The efferent arteriole is contracted so that the intraglomerular pressure increases and thus filtration and the GFR increase too

000015def24e00ca30

64

A 75-year-old man is on haemodialysis. The dialysis catheter is located in the jugular vein. He attends the Emergency department because he is short of breath, lacks energy and has a temperature. At examination his blood pressure is 90/65, there is slight tenderness around the catheter opening, but not redness or pus.

What condition does he most probably have?

- A** Septicaemia
- B** Overhydration
- C** Pneumocystis jirovecii pneumonia
- D** Bronchitis

000015def24e00ca30

65

A 35-year old woman has biopsy-verified IgA nephritis. She is receiving treatment with ACE inhibitors for hypertension and reduction of proteinuria.

There is slow disease progression. At the last check-up at the Kidney Outpatient Clinic, her creatinine was 300 micromol/L, eGFR 18 ml/min, and potassium 5.0 mmol/L.

She is more tired than previously, her weight is stable and she is still in a full-time job.

What would you do now?

- A Start the evaluation process for a kidney transplant.
- B Refer her for AV fistula or PD catheter
- C Start dialysis immediately
- D Continue with the same treatment and give the patient an appointment for check-up in 4 months.

000015de624e00ca30

66

Nephrotic syndrome is characterised by general oedema and significant changes in blood test results. Which of the following combinations of blood test results belong most probably to such a patient?

(ref. values: Hb 11,7 - 15,3 g/dL (female), 13,4 - 17,0 g/dL (male), s-creatinine 45 - 90 µmol/L (female), 60 - 105 µmol/L (male), s-cholesterol >=50 years: 3,9 - 7,8 mmol/L, s-albumin 40-69 years: 36-45 g/L)

- A Hb 13.2 g/dl, s-creatinine 185 umol/L, s-cholesterol 6.9 mmol/L, s-albumin 28 g/L
- B Hb 11.2 g/dl, s-creatinine 85 umol/L, s-cholesterol 5.9 mmol/L, s-albumin 33 g/L
- C Hb 11.2 g/dl, s-creatinine 85 umol/L, s-cholesterol 12.2 mmol/L, s-albumin 23 g/L
- D Hb 11.2 g/dl, s-creatinine 85 umol/L, s-cholesterol 5.9 mmol/L, s-albumin 62 g/L

000015de624e00ca30

67

At the health centre you see a 47-year old man who has previously been healthy. He feels unwell and has a headache, and thinks his right arm feels a little numb. He does not smoke, takes no medication and has a BMI of 28 (BMI 25.0-29.9 overweight).

Lab: BP 228/140. Creatinine 90 µmol/L (60-105 µmol/L), eGFR 62 mL/min/1.73 m² (>60 mL/min/1.73m²), Urine dipstick: ++albumin.

Which of the following assessments is most correct?

- A You admit the patient to the Dept. of Medicine for emergency help.
- B You order rush CT of the head and give the patient an appointment for follow-up the next day if there are no findings from the investigation.
- C You start treatment with a calcium-antagonist and give him an appointment for follow-up in 3 days' time.
- D You start blood pressure treatment with an angiotensin receptor-blocker and give the patient a new appointment in 2 days' time.

000015de624e00ca30

68

When considering starting treatment for hypertension, it is useful to have information on any subclinical organ damage.

Which of the following test combinations has the best benefit/cost ratio?

- A s-creatinine, s-uric acid, echocardiography
- B u-dipstick, s-creatinine, ECG
- C s-creatinine, ophthalmoscopy, echocardiography
- D ECG, echocardiography, ultrasound carotid vessels

000015de624e00ca30

69

The definition of hypertension is a systolic blood pressure above 140 or diastolic blood pressure above 90 mmHg measured several times under standard conditions at the doctor's surgery. The treatment target is <140/90 mmHg for most patients, but there are exceptions.

Which of the groups below should have a lower target level after treatment?

- A** Patients with hypertensive nephrosclerosis (arterionephrosclerosis).
 - B** Patients with IgA glomerulonephritis.
 - C** Patients with impaired renal function after a heart attack.
 - D** Patients with chronic renal disease and proteinuria demonstrated by u-dipstick.
-

000015de24e00ca30

70

Anna (82 years old) has COPD, diabetes mellitus type II and chronic renal failure and must start on renal replacement therapy. She is overweight (BMI 30) and has previously undergone several abdominal operations. The nearest haemodialysis unit is 1.5 hours away, each way.

Which method of dialysis is best for her and why?

- A** Haemodialysis at the nearest hospital because she is not suitable for peritoneal dialysis.
 - B** Haemodialysis at the nearest hospital because peritoneal dialysis is too demanding for the municipal home nursing services.
 - C** Haemodialysis at the nearest hospital because peritoneal dialysis is contraindicated in diabetes mellitus.
 - D** Peritoneal dialysis in the home due to the long travel time.
-

000015de24e00ca30

71

A 69-year old woman has measured her blood sugar in capillary blood using her sister's equipment for measuring blood sugar at home. This gave a result of 13.4 mmol/L about an hour after she had drunk coffee and eaten apple cake with cream. She feels well. Her doctor performed a glucose tolerance test which gave the following results for p-glucose:

- Fasting sample (0 sample): 7.9 mmol/L
- Sample 2h after glucose intake: 9.9 mmol/L

Reference range for p-glucose is 4.2-6.3 mmol/L

What is the most correct classification of the patient based on the above information?

- A** Diabetes mellitus
 - B** Both diabetes mellitus and impaired glucose tolerance
 - C** Impaired glucose tolerance
 - D** Neither diabetes mellitus nor impaired glucose tolerance
-

000015de24e00ca30

72

A 69-year old woman went to her GP because she had felt tired and lacked energy more than usual over the last 6 months. A fasting blood sample was taken at 8.00 am the results of which are shown below.

Analysis	Unit	Reference range	Result (Flag)
S-Glucose	mmol/L	4.0 - 6.0	6.5 (H)
B-HbA1c (glycated haemoglobin)	mmol/mol	28 - 40	44 (H)
S-Calcium, total	mmol/L	2.15 - 2.51	2.53 (H)
S-Calcium, free in serum (standardised at pH 7.40)	mmol/L	1.18 - 1.32	1.29
S-Cortisol	nmol/L	At: 06-10: 133 - 537 At: 16-20: 68 - 327	343
S-Anti-thyroid peroxidase (TPO)	kU/L	< 35	111 (H)
S-Phosphate	mmol/L	0.85 - 1.50	1.12
S-Free T4	pmol/L	12.0 - 22.0	8.6 (L)
S-PTH (Parathyroid hormone)	pmol/L	1.6 – 6.9	6.0
S-TSH (Thyroid stimulating hormone)	mIU/L	0.27 – 4.20	18 (H)
Anti-TSH receptor (TRAS)	IU/L	<1.0	0.1

Based on the medical history and the results of the biochemical analyses, what is the most probable diagnosis?

- A** Adrenal cortex insufficiency
- B** Primary hypothyroidism due to autoimmune thyroiditis
- C** Diabetes mellitus type II
- D** Hypercalcemia due to hyperparathyroidism

000015de624e00ca30

73

Which two classes of antidiabetics should primarily be chosen in cases of known heart disease when metformin and diet/physical activity do not provide adequate glucose regulation?

- A** DPP4 inhibitor or insulin
- B** SGLT2 inhibitor or GLP1 analogue
- C** Sulphonylurea or SGLT2 inhibitor
- D** DPP4 inhibitor or GLP1 analogue

000015de624e00ca30

74

A 60-year old woman comes to see you because she has had episodes of sweating that have increased over the last months. Combined with this she has also had palpitations and anxiety. You measure her blood pressure which is considerably raised. Some years ago the patient was diagnosed with primary hyperparathyroidism and was treated surgically. You refer her now for CT of the adrenals which reveals a left-sided tumour. What is the most probable diagnosis?

- A** Conn's syndrome (aldosterone-producing tumour)
- B** Acromegaly
- C** Pheochromocytoma as part of multiple endocrine neoplasia (MEN) type 2
- D** Cushing's syndrome

000015de624e00ca30

75

A man aged 55 has been shown to have hypercalcemia and elevated PTH compatible with primary hyperparathyroidism. He has normal kidney function. Which of these statements can give a corresponding pattern of blood test results?

- A** tertiary hyperparathyroidism
- B** familial hypocalciuric hypercalcemia
- C** secondary hyperparathyroidism
- D** myelomatosis

000015del24e00ca30

76

A 35-year old woman was diagnosed with autoimmune hyperthyroidism (Graves' disease) 1 year ago. She is being treated with an anti-thyroid drug (carbimazole). Over the last six months, metabolic results have been normal with free thyroxine (FT4) in the reference range (11.6-19.1 pmol/L) and thyroid stimulating hormone (TSH) around 1.0 mIU/L (ref. 0.24-3.78 mIU/L). The last month she has been tired, slightly nauseous and dizzy, and has lost about 4-5 kg in weight. Regular periods. Blood pressure 90/60. Normal kidney function, serum sodium 136 mmol/L (ref. 137-145 mmol/L) and potassium 4.3 mmol/L (ref. 3.6-4.6 mmol/L). FT4 is now 15.6 pmol/L and TSH is 0.97 mIU/L. Which of the blood test results below is the most important for clarifying the condition?

- A** TSH receptor antibody (TRAS) and anti-TPO (anti-thyroid peroxidase)
- B** Follicle stimulating hormone (FSH) and Luteinizing hormone (LH)
- C** Prolactin
- D** Cortisol and ACTH

000015del24e00ca30

77

A 51-year old woman has hyperthyroidism with free thyroxine (FT4) 51.2 pmol/L (reference range 11.6-19.1 pmol/L) and thyroid stimulating hormone (TSH) <0.01 mIU/L (ref. 0.24-3.78 mIU/L). TSH receptor antibody is 12.4 (ref. <1.5) IU/L. Moderate endocrine ophthalmopathy. Two weeks after starting on an anti-thyroid drug (carbimazole) she has an itchy exanthem on her trunk. What do you recommend she should do?

- A** That she discontinues carbimazole and takes new blood tests in two weeks
- B** That she discontinues the carbimazole and is treated with radioactive iodine (RAI)
- C** That she switches to a different anti-thyroid drug (propylthiouracil=PTU) and take an antihistamine.
- D** That she discontinues carbimazole and is treated surgically

000015del24e00ca30

78

A woman, 61 years of age, has newly diagnosed type 2 diabetes. She complains of numbness in her feet and wonders if this can have anything to do with her diabetes. Which examination would you do first considering that this could be diabetic polyneuropathy (DPN)?

- A** 10g monofilament test
- B** Palpate the foot pulses
- C** Examine the patella reflex
- D** Refer for neurography

000015del24e00ca30

79

A 78-year old woman with multimorbidity lives in a care home and has had type 1 diabetes for 36 years. She uses insulin multi-injections and has problems with repeated hypoglycemic episodes with confusion and anxiousness. What is the treatment target (HbA1c and/or blood glucose) for this patient?

- A** HbA1c between 53-64 mmol/mol (7.0-8.0%). Fasting blood glucose 5-8 mmol/L
- B** That the patient avoids symptomatic hyperglycemia, i.e. the blood glucose rarely exceeds 12-14 mmol/L
- C** HbA1c about 53 mmol/mol (7%). Fasting blood glucose 4-7 mmol/L
- D** HbA1c about 48 mmol/mol (6.5%). Fasting blood glucose 4-6 mmol/L

000015del24e00ca30

80

A 42-year old man has had type 1 diabetes (T1D) for 25 years. He has had laser treatment on both eyes due to diabetic retinopathy and now has a corrected visual acuity of 0.4+ on the right eye and 0.4+ on the left eye. Normal field of vision. He wants a doctor's certificate for a driving licence for private cars (group 1). Does he meet the health requirements for a driving licence for group 1 in regard to visual acuity?

- A** He meets the health requirements in regard to visual acuity
 - B** It is unclear whether he meets the health requirements in regard to visual acuity
 - C** He does not meet the health requirements in regard to visual acuity
-

000015de124e00ca30

81

A 55-year old woman has lost weight, has palpitations, slight tremors and signs of periorbital oedema. Moderately enlarged thyroid gland with even consistency, no nodules, no palpation tenderness. Free thyroxine (FT4) is 31.8 (reference range 11.6-19.1 pmol/L) and thyroid stimulating hormone (TSH) <0.01 mIU/L (ref. 0.24-3.78 mIU/L). Which additional blood tests or investigations should be done first to clarify whether she has autoimmune hyperthyroidism (Graves' disease)?

- A** Thyroid scintigraphy
 - B** C-reactive protein (CRP)
 - C** TSH receptor antibody
 - D** Anti-TPO = anti-thyroid peroxidase antibody
-

000015de124e00ca30

82

A woman (aged 64) has felt increasingly tired and worn out over the last 1-2 years. She takes no regular medicines. She goes to her GP who measures her BP 115/65 and pulse 56. Initial blood tests show TSH 2.4 mIU/L (0.24-4.2) and free T4 7.6 pmol/L (11.6-19.1).

Which of the following laboratory tests are most relevant in further differential diagnosis?

- A** Anti-TPO
 - B** Salivary cortisol levels in the morning and evening
 - C** Free T3
 - D** FSH, LH, prolactin and cortisol in serum (morning)
-

000015de124e00ca30

83

A 58-year old woman sees you as her GP about problems with one of her breasts. The last few weeks, her left breast has been slightly more tender than her right, but not acutely painful. The woman is otherwise healthy and takes no medicines. She has never breastfed. At examination you see the following skin changes on her breast:



You palpate both breasts and find a 3 cm tumour in the upper lateral quadrant of the left breast. You do not palpate clearly enlarged lymph nodes on either side. Which statement is the most correct?

- A** The skin changes indicate breast cancer with metastases, and the woman must be referred for mammography
- B** The skin changes indicate a fibroadenoma in the breast, but the woman should nonetheless be referred to the Diagnostic Breast Centre.
- C** The skin changes indicate locally advanced breast cancer, and the woman must be referred to the Diagnostic Breast Centre
- D** The skin changes indicate mastitis, but the woman should still be referred for mammography

000015de624e00ca30

84

In your office you have a 62-year old woman with primary hyperparathyroidism. The condition has just recently been diagnosed, and treatment has not yet started. When asked about symptoms, the patient reports having been very thirsty with frequent toilet visits, tingling in fingers and toes, cramps in both legs and muscle weakness. She feels lethargic and is often constipated.

Which of the symptoms mentioned fit the diagnosis of primary hyperparathyroidism well, and why?

- A** Tingling in fingers and toes, muscle weakness and lethargy due to hypocalcemia
- B** Increased thirst, muscle weakness and constipation due to hypocalcemia
- C** Increased thirst, muscle weakness and constipation due to hypercalcemia
- D** Tingling in fingers and toes, muscle weakness and lethargy due to hypercalcemia

000015de624e00ca30

85

A 55-year old woman sees you as her GP after she has noticed several episodes with secretion from her left breast.

Should this be investigated further?

- A** Secretion from the breast does not need to be investigated further unless one finds a tumour at clinical examination.
 - B** Secretion from the breast should always be investigated, and after taking a medical history and performing a clinical examination, you refer her.
 - C** If the medical history reveals clear or bloody fluid from her breast, the patient must be referred for investigation for a tumour in the milk duct.
 - D** If the medical history reveals clear or bloody fluid from her breast, the patient must be referred for investigation for breast cancer.
-

000015def24e00ca30

86

A 60-year old man has an appointment with his GP because of a palpable tumour in his thyroid. He has no discomfort from his throat, but discovered the tumor when he was looking in the mirror while shaving. The tumour is well-defined, appears to be movable and has a relatively soft consistency. It feels like it is about 3cm at its largest. His basal metabolism is normal.

How should this be investigated further?

- A** The patient should be referred for CT of the neck
 - B** The patient should be referred to an endocrinological outpatient clinic for investigation including an ultrasound scan and maybe a fine needle biopsy.
 - C** Further investigation is not necessary as he has no discomfort.
 - D** Thyroid scintigraphy should be performed. If cold nodules are found these should be removed surgically.
-

000015def24e00ca30

87

A 66-year old man has been shown to have an elevated serum PSA by his GP, and has been investigated further through the Cancer Care Pathway for prostate cancer. Biopsy reveals prostate cancer with Grade Group 4 (Gleason 4+4). The Urologist is considering radical prostatectomy. Which imaging diagnostic is best suited to staging to take into consideration metastases?

- A** Metastasis investigation is not necessary in Grade Group 4.
 - B** MRI prostate.
 - C** CT of lungs (thorax) and abdomen.
 - D** MRI pelvis and spine
-

000015def24e00ca30

88

A 74-year old man has hypertension (180/100mmHg) and is being treated for diabetes mellitus with metformin tablets. He is now being investigated for glomerulonephritis because of proteinuria and oedema. You want an ultra-sound guided kidney biopsy. Which preparation of the patient is important to consider in this situation?

- A** Ensure good hydration; if necessary give fluids intravenously.
 - B** Get his blood pressure under control with antihypertensives.
 - C** Discontinue metformin.
 - D** Give prophylactic antibiotics.
-

000015def24e00ca30

89

A 23-year old woman has had a lot of problems with recurring urinary tract infections. She has now been admitted to hospital with symptoms of infection in the upper urinary tract, with a high fever in spite of antibiotic treatment for 3 days through her GP. Which imaging diagnostic should be the first choice?

- A CT urinary tract with i.v. contrast in the parenchymal phase.
- B Ultrasound scan of the urinary tract.
- C 3-phase CT of the urinary tract.
- D Stone-CT urinary tract without i.v. contrast.

000015def24e00ca30

90

What is the most important limitation of ultrasound in regard to concrements when investigating the urinary tract?

- A Ultrasound does not provide images of chalk (calcium carbonate)-containing lesions.
- B Ultrasound cannot distinguish between a kidney stone, gallstone or arterial chalk (calcium carbonate).
- C Peristalsis in the ureter wall gives movement artefacts.
- D Intestinal gas artefacts hinder inspection of the ureters.

000015def24e00ca30

91

In connection with hospitalisation for left flank pain, CT without i.v. contrast was performed which showed an 8 mm concrement in the left ureteropelvic transition. Eventually, the patient's pain was alleviated and the kidney was relieved using a JJ-stent. The patient was discharged feeling well the next day. The patient was given an Outpatient appointment the following week when treatment with ESWL was planned. They wanted to confirm whether the concrement was still present and whether it was in the same location as previously. Which imaging modality is usually used to check the location of a urinary tract concrement?

- A CT urinary tract with i.v. contrast, in 3 phases.
- B Stone-CT urinary tract without i.v. contrast.
- C X-ray urinary tract.
- D Ultrasound of the urinary tract.

000015def24e00ca30

92

When using CT to decide whether a demonstrated kidney lesion is a solid, malignancy-suspect tumour, a combination of several contrast phases is preferably used. How are the phases normally combined in this instance, and why?

- A Precontrast phase and parenchymal phase. An increase in density between the phases verifies that it is a circumscribed tumour.
- B Precontrast phase and arterial phase. Verifies calcifications in the tumour and the vessels that supply the tumour.
- C Precontrast phase, parenchymal phase and excretory phase. Can measure the density in the phases and calculate the absolute and relative washout.
- D Arterial phase and parenchymal phase. Visualises both vessel supply and the extent of the tumour.

000015def24e00ca30

93

Vesicoureteral reflux (VUR) is a condition in which the urine flows back from the bladder and up into the ureters, and eventually to the renal pelvis. VUR is assumed to occur in 1-2% of all children. Reflux is demonstrated in 15-50% of children with urinary tract infections. Which imaging modality is used to detect and grade VUR?

- A CT urography (with intravenous contrast)
- B X-ray - MCUG
- C Diuresis renography
- D Ultrasound without contrast

000015def24e00ca30

94

Wet lungs (TTN/Transient Tachypnea of the Newborn) is a benign condition that is due to the retention of a small amount of amniotic fluid in the lung tissue after birth. This is the most common cause of respiratory problems in the newborn. The condition manifests as mild breathing problems immediately, or within a few hours, after the birth. Symptoms can persist for a few hours to a few days (max. 48-72 hours).

Which of the following alternatives is a well-known risk factor for wet lungs?

- A Discoloured amniotic fluid
- B Renal disease in the mother
- C Caesarian section
- D Extreme prematurity (<28 weeks)

000015d0f24e00ca30

95

A 40-year old woman has recently undergone surgery for breast cancer. She tells you that the tumour is human epidermal growth factor receptor 2 (HER2) positive. She asks you a number of questions about this.


Which statement is correct?

- A HER2-positive breast cancer tumours are HER2-amplified
- B HER2-positive breast cancer tumours have little HER2 protein
- C HER2-positive breast cancer tumours are BRCA1-mutated
- D HER2-positive breast cancer tumours are often low-grade

000015d0f24e00ca30

96

A 34-year old woman undergoes surgical resection of a tumour-suspected area in the upper, lateral quadrant, right breast. The image shows a hematoxylin and eosin (H&E) stained histology section from the area (x100).




Which diagnosis best fits the findings in the image?

- A Ductal carcinoma in situ
- B Metastasis from malignant melanoma
- C Infiltrating carcinoma
- D Normal breast tissue

000015del24e00ca30

97

A 24-year old woman has had an enlarged thyroid for several months. It started during pregnancy and the thyroid was tender during the first weeks. Now she has symptoms of hypothyroidism. The image shows a hematoxylin and eosin (H&E) stained histology section from her thyroid (x600).



What is the diagnosis?

- A Granulomatous thyroiditis
- B Papillary carcinoma
- C Lymphocytic thyroiditis (Hashimoto's disease)
- D Hyperplasia (Grave's disease)

000015del24e00ca30

98

One of your patients was operated with removal of a tumour in her left ovary. You receive the pathology report. The report states that the tumour contains respiratory tract epithelium, squamous epithelium, sebaceous glands and hair follicles.

Based on the findings in the pathology report, which statement is correct?

- A The tumour is probably a metastasis from a skin tumour
- B The tumour is probably a serous ovarian cancer
- C The tumour is probably a stromal (sex cord stromal) tumour
- D The tumour is probably a germinal cell tumour

000015del24e00ca30

99

A 60-year old woman comes to you because of postmenopausal bleeding. After investigation by taking a pipelle sample she is diagnosed with endometrial hyperplasia with atypia.

Which statement about endometrial hyperplasia is correct?

- A Endometrial hyperplasia with atypia does not need treatment, but the woman should be followed-up every fifth year.
- B Endometrial hyperplasia with atypia means that there is an infiltrating growth
- C One generally sees atrophic glands and little stroma in endometrial hyperplasia.
- D Endometrial hyperplasia with and without atypia have different risks of developing cancer

000015del24e00ca30

100

A 38-year old man has recently undergone surgery for removal of a testicular tumour. The tumour has been diagnosed to be a seminoma.


Which statement is correct?


- A Seminomas are classified as germinal cell tumours
- B Seminomas occur most frequently in men aged between 70 and 80
- C Stromal (sex cord stromal) tumours are the most common tumour in the testis
- D Most testicular tumours grow rapidly and cause pain

000015del24e00ca30

101

A 76-year old man has had pain in his lumbar region for quite some time, and occasionally has noticed haematuria. Ultrasound scan reveals a 5 cm diameter, almost round tumour in the upper pole of the left kidney. The images show histopathological sections from the tumour (H&E, 100x and 400x).





What is the most probable diagnosis?


- A** Ectopic adrenal cortex
- B** Wilms' tumour
- C** Lipoma
- D** Renal cell carcinoma

000015d0e24e00ca30

102

A 60-year old man was referred for cystoscopy due to persistent haematuria. A biopsy was taken from a lesion in the urinary bladder mucous membrane. The image shows a section from this area (H&E stain, x100).

What is the diagnosis?



- A** Invasive urothelial carcinoma, WHO grade 3
- B** Granulomatous cystitis
- C** Squamous cell metaplasia
- D** Papillary urothelial carcinoma, WHO grade 1

000015del24e00ca30

103

Growth of more than one bacterial strain in a urine sample is often interpreted as a sign of contamination at sampling.

In which cases do this phenomenon most probably represent a true mixed infection?

- A** When the sample is collected as a mid-stream sample
- B** In cases of known anomaly of the urinary tract
- C** In men with prostatic hypertrophy
- D** When the sample is collected using a single-use urinary catheter

000015del24e00ca30

104

In a six-month old child with a fever, a urine sample collected using a plastic bag sample reveals 10^4 CFU/ml of *E. coli*.

How should this finding be best interpreted in regard to the probable cause of urinary tract infection?

- A** The finding cannot be taken into account because the plastic bag sample is best suited for gram-positive bacteria that are not normally present in the perineum
- B** As the requirement for significant bacteriuria in infants has been set at 10^4 CFU/ml, the finding must be assumed to represent the probable cause of the urinary tract infection
- C** The finding is not significant because the use of a urinary plastic bag sample requires a bacterial count of at least 10^5 CFU/ml
- D** If other causes of the fever cannot be found, the finding must be presumed to represent the probable cause of the urinary tract infection

000015del24e00ca30

105

The term primary urinary tract pathogenic bacterial strains is used about bacterial strains that can cause urinary tract infections in people with normal urinary tracts. Which bacterial strain belongs to this group?

- A Chlamydia trachomatis
 - B Staphylococcus saprophyticus
 - C Enterococcus faecalis
 - D Klebsiella oxytoca
-

000015d0e24e00ca30

106

In your practice, you get the results from the urinary culture of a patient with pyelonephritis. The clean culture gives E. coli 10^5 CFU/ml which is sensitive to nitrofurantoin, gentamicin, trimethoprim sulfa and cefotaxime.

Which of these medications is the best suited to treat this patient in general practice?

- A Nitrofurantoin
 - B Cefotaxime
 - C Trimethoprim sulfa
 - D Gentamicin
-

000015d0e24e00ca30

107

A young woman has just returned from holiday in Tanzania and attends your office with a fever of 39.5 degrees Celsius. She has a urticarial rash on her thighs and stomach. She hasn't noticed that she has been bitten by an insect, but says that this started a few days after she went swimming in Lake Victoria, a freshwater lake.

What is the most probable diagnosis?

- A Acute schistosomiasis (Katayama fever)
 - B Leishmaniasis (Kala-azar)
 - C Rickettsiosis
 - D Typhoid fever
-

000015d0e24e00ca30

108

Patients with rheumatoid arthritis are increasingly taking tocilizumab (RoActemra®) as immunosuppressive treatment. This medicine blocks the Interleukin-6 receptor.

When diagnosing infections, which problems are associated with this medication?

- A Reduced pain response and absent leukocytosis
 - B Reduced fever response and absent CRP increase
 - C Reduced pain response and absent CRP increase
 - D Reduced experience of dyspnoea and absent leukocytosis
-

000015d0e24e00ca30

109

A 20-year old woman has had unprotected sex (vaginal intercourse) with a man who has subsequently been shown to be HIV positive although he was unaware of this at the time of intercourse. What are the chances that she is infected?

- A 5-10%
 - B 2-5%
 - C Less than 1%
 - D More than 15%
-

000015d0e24e00ca30

110

Treatment for a urinary tract infection has been started in an 85-year old woman with a fever of 38.5°C admitted to a care home. Urine culture reveals clean culture of *Proteus mirabilis* (belongs to the Enterobacteriaceae family) with 10⁴ CFU/ml, with the following sensitivity profile:

ampicillin R
cefuroxime I
nitrofurantoin S
mecillinam S
trimethoprim S

Which of the following medicines is best suited to treatment of this patient in the care home?

- A Trimethoprim
- B Cefuroxime
- C Mecillinam
- D Nitrofurantoin
- E Ampicillin

000015de624e00ca30

111

Angiotensin receptor antagonists are often used for hypertension and heart failure.

Which of the following conditions warrants extra monitoring of the patient after starting on an angiotensin receptor antagonist?

- A Aortic insufficiency
- B Intermittent claudication
- C Prior tendency to syncope when treated with nitroglycerine
- D Tendency for hyperkalaemia

000015de624e00ca30

112

Alpha-blockers have a vasodilatory effect and are marketed, among other indications, as blood pressure lowering agents.

Why is this group of medicines not used extensively in the treatment of high blood pressure?

- A It has a diabetogenic effect
- B It can cause severe hypokalaemia
- C It can cause impotence
- D It increases the risk of heart failure

000015de624e00ca30

113

A patient has newly-diagnosed osteoporosis and you are considering treatment with alendronate and calcium. An interaction search displays a warning symbol and the following text:

M05B A Bisphosphonates - A12A A Calcium

Decreased absorption of oral bisphosphonates.

What is the most correct handling of this interaction?

- A You increase the dose of alendronate by 50%
- B Alendronate and calcium cannot be combined
- C Ask the patient to take alendronate before breakfast and calcium in the evening
- D You ask the patient to take alendronate and calcium every other day

000015de624e00ca30

114

Contraceptive agents contain one or more hormones.

Which of the following contraceptive agents does *not* contain oestrogen?

- A NuvaRing contraceptive ring
- B Nexplanon contraceptive rod implant
- C Evra contraceptive patches
- D Loette contraceptive pills

000015de624e00ca30

115

A 72-year old woman with diabetes type 2, elevated blood pressure and hypercholesterolaemia has taken the following medicines over the last six months:

- Metformin (a biguanide derivative, for diabetes)
- Empagliflozin (a sodium-glucose-cotransporter-2 inhibitor (SGLT2 inhibitor), for diabetes)
- Valsartan (an angiotensin II receptor antagonist, for high blood pressure)
- Atorvastatin (a statin, for elevated cholesterol)

Over the last months the patient has had very frequent, recurring urinary tract infections.

Based on this, which of the patient's medicines should you consider discontinuing?

- A Atorvastatin
- B Empagliflozin
- C Valsartan
- D Metformin

000015de624e00ca30

116

After the American "Women's Health Initiative" study on postmenopausal oestrogen therapy was published in 2002, there was a significant drop in this treatment in a number of industrialised countries. In the wake of this reduction in postmenopausal oestrogen treatment, epidemiologists could also measure a reduction in a certain type of cancer in postmenopausal women.

Which type of cancer showed a drop in incidence?

- A Colon cancer
- B Endometrial cancer
- C Vaginal cancer
- D Breast cancer

000015de624e00ca30

117

In connection with major trauma, some patients may suffer a large, but often reversible, reduction in renal function. Such renal function impairment may result in the contraindication of central groups of analgesics. Among the most used analgesic medications we find the opioids morphine and codeine, the NSAID diclofenac, and paracetamol.

One of these four drugs may be used at recommended doses without taking into account the patient's kidney function. Which one?

- A Codeine
- B Diclofenac
- C Morphine
- D Paracetamol

000015de624e00ca30

118

A woman is solely breastfeeding her two-month old child. She must take a medicine which potentially could have an effect on the breastfed child. To evaluate how safe the medicine would be for the child, you must calculate the child's *theoretical dose*, i.e. how much medicine the child theoretically will take up.

How is the theoretical dose calculated?

- A The amount of milk consumed (about 150 ml per kg bodyweight per day) is multiplied by the concentration of the medicine in the mother's serum
- B The serum concentration of the medicine is measured in the child's blood and multiplied by the child's blood volume (about 500 ml)
- C The amount of milk consumed (about 150 ml per kg bodyweight per day) is multiplied by the concentration of the medicine in the milk
- D The amount of milk consumed (about 150 ml per kg bodyweight per day) is divided by the concentration of the medicine in the milk

000015de624e00ca30

119

A young woman takes lamotrigine as indicated for epilepsy. She is well-regulated on a relatively low dose. She is planning to become pregnant.

What should you tell her?

- A** She should discontinue lamotrigine before she becomes pregnant
 - B** She should be monitored with repeated serum concentration measurements and any necessary dose adjustments before, during and after the pregnancy.
 - C** She should switch from lamotrigine to valproate before she becomes pregnant
 - D** It is unlikely that the pregnancy will affect the dose requirements, and monitoring other than routine is not necessary
-

000015de624e00ca30

120

A 75-year old man takes isosorbide dinitrate as indicated for angina pectoris. He sees you as his GP because he wants a prescription for sildenafil for erectile dysfunction. An interaction search reveals possible interaction between isosorbide dinitrate and sildenafil.

What undesired effect can occur if these two medicines are combined?

- A** Sildenafil induces metabolism of isosorbide dinitrate and gives an increased risk of therapy failure
 - B** The combination will increase the blood pressure lowering effect and give an increased risk of hypotension
 - C** The combination will lower the blood pressure reducing effect and will give an increased risk of angina pectoris symptoms
 - D** Sildenafil inhibits metabolism of isosorbide dinitrate and gives an increased risk of overdosing
-

000015de624e00ca30